

# 3D Computer Vision

Radim Šára    Martin Matoušek

Center for Machine Perception  
Department of Cybernetics  
Faculty of Electrical Engineering  
Czech Technical University in Prague

<https://cw.fel.cvut.cz/wiki/courses/tdv/start>

<http://cmp.felk.cvut.cz>

<mailto:sara@cmp.felk.cvut.cz>

phone ext. 7203

rev. September 22, 2020



Open Informatics Master's Course

# Module II

## Perspective Camera

- 2.1 Basic Entities: Points, Lines
- 2.2 Homography: Mapping Acting on Points and Lines
- 2.3 Canonical Perspective Camera
- 2.4 Changing the Outer and Inner Reference Frames
- 2.5 Projection Matrix Decomposition
- 2.6 Anatomy of Linear Perspective Camera
- 2.7 Vanishing Points and Lines

**covered by**

[H&Z] Secs: 2.1, 2.2, 3.1, 6.1, 6.2, 8.6, 2.5, Example: 2.19

## ► Basic Geometric Entities, their Representation, and Notation

- entities have names and representations
- names and their components:

entity	in 2-space	in 3-space
point	$m = (u, v)$	$X = (x, y, z)$
line	$n$	$O$
plane		$\pi, \varphi$

- associated vector representations

$$\mathbf{m} = \begin{bmatrix} u \\ v \end{bmatrix} = [u, v]^{\top}, \quad \mathbf{X} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, \quad \mathbf{n}$$

will also be written in an 'in-line' form as  $\mathbf{m} = (u, v)$ ,  $\mathbf{X} = (x, y, z)$ , etc.

- vectors are always meant to be columns  $\mathbf{x} \in \mathbb{R}^{n \times 1}$
- associated homogeneous representations

$$\underline{\mathbf{m}} = [m_1, m_2, m_3]^{\top}, \quad \underline{\mathbf{X}} = [x_1, x_2, x_3, x_4]^{\top}, \quad \underline{\mathbf{n}}$$

'in-line' forms:  $\underline{\mathbf{m}} = (m_1, m_2, m_3)$ ,  $\underline{\mathbf{X}} = (x_1, x_2, x_3, x_4)$ , etc.

- matrices are  $\mathbf{Q} \in \mathbb{R}^{m \times n}$ , linear map of a  $\mathbb{R}^{n \times 1}$  vector is  $\mathbf{y} = \mathbf{Q}\mathbf{x}$
- $j$ -th element of vector  $\mathbf{m}_i$  is  $(\mathbf{m}_i)_j$ ; element  $i, j$  of matrix  $\mathbf{P}$  is  $\mathbf{P}_{ij}$

## ► Image Line (in 2D)

a finite line in the 2D  $(u, v)$  plane

$$a u + b v + c = 0$$

has a parameter (homogeneous) vector

$$\underline{\mathbf{n}} \simeq (a, b, c), \quad \|\underline{\mathbf{n}}\| \neq 0$$

and there is an equivalence class for  $\lambda \in \mathbb{R}, \lambda \neq 0$   $(\lambda a, \lambda b, \lambda c) \simeq (a, b, c)$

### 'Finite' lines

- standard representative for finite  $\underline{\mathbf{n}} = (n_1, n_2, n_3)$  is  $\lambda \underline{\mathbf{n}}$ , where  $\lambda = \frac{1}{\sqrt{n_1^2 + n_2^2}}$   
assuming  $n_1^2 + n_2^2 \neq 0$ ;  $\mathbf{1}$  is the unit, usually  $\mathbf{1} = 1$

### 'Infinite' line

- we augment the set of lines for a special entity called the **line at infinity** (ideal line)

$$\underline{\mathbf{n}}_\infty \simeq (0, 0, 1) \quad (\text{standard representative})$$

- the set of equivalence classes of vectors in  $\mathbb{R}^3 \setminus (0, 0, 0)$  forms the projective space  $\mathbb{P}^2$   
a set of rays  $\rightarrow 21$
- line at infinity is a proper member of  $\mathbb{P}^2$
- I may sometimes wrongly use  $=$  instead of  $\simeq$ , if you are in doubt, ask me

## ► Image Point

Finite point  $\mathbf{m} = (u, v)$  is incident on a finite line  $\mathbf{n} = (a, b, c)$  iff      iff = works either way!

$$a u + b v + c = 0$$

can be rewritten as (with scalar product):  $(u, v, \mathbf{1}) \cdot (a, b, c) = \mathbf{m}^\top \mathbf{n} = 0$

### 'Finite' points

- a finite point is also represented by a homogeneous vector  $\mathbf{m} \simeq (u, v, \mathbf{1})$ ,  $\|\mathbf{m}\| \neq 0$
- the equivalence class for  $\lambda \in \mathbb{R}, \lambda \neq 0$  is  $(m_1, m_2, m_3) = \lambda \mathbf{m} \simeq \mathbf{m}$
- the standard representative for finite point  $\mathbf{m}$  is  $\lambda \mathbf{m}$ , where  $\lambda = \frac{1}{m_3}$  assuming  $m_3 \neq 0$
- when  $\mathbf{1} = 1$  then units are pixels and  $\lambda \mathbf{m} = (u, v, 1)$
- when  $\mathbf{1} = f$  then all elements have a similar magnitude,  $f \sim$  image diagonal

use  $\mathbf{1} = 1$  unless you know what you are doing;

all entities participating in a formula must be expressed in the same units


### 'Infinite' points

- we augment for **points at infinity** (ideal points)  $\mathbf{m}_\infty \simeq (m_1, m_2, 0)$   
proper members of  $\mathbb{P}^2$
- all such points lie on the line at infinity (ideal line)  $\mathbf{n}_\infty \simeq (0, 0, 1)$ , i.e.  $\mathbf{m}_\infty^\top \mathbf{n}_\infty = 0$

## ► Line Intersection and Point Join

The point of **intersection**  $m$  of image lines  $n$  and  $n'$ ,  $n \neq n'$  is

$$\underline{m} \simeq \underline{n} \times \underline{n}'$$




**proof:** If  $\underline{m} = \underline{n} \times \underline{n}'$  is the intersection point, it must be incident on both lines. Indeed, using known equivalences from vector algebra

$$\underline{n}^\top \underbrace{(\underline{n} \times \underline{n}')}_{\underline{m}} \equiv \underline{n}'^\top \underbrace{(\underline{n} \times \underline{n}')}_{\underline{m}} \equiv 0$$

The **join**  $n$  of two image points  $m$  and  $m'$ ,  $m \neq m'$  is

$$\underline{n} \simeq \underline{m} \times \underline{m}'$$



Parallel lines intersect (somewhere) on the line at infinity  $\underline{n}_\infty \simeq (0, 0, 1)$ :

$$a u + b v + c = 0,$$

$$a u + b v + d = 0,$$

$$d \neq c$$

$$(a, b, c) \times (a, b, d) \simeq (b, -a, 0)$$

- all such intersections lie on  $\underline{n}_\infty$
- line at infinity therefore represents the set of (unoriented) directions in the plane
- Matlab: `m = cross(n, n_prime);`

Thank You